

MAIN PANEL

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Chipco[®] Baycor[®]

FUNGICIDE

ACTIVE CONSTITUENT: 300 g/L BITERTANOL
SOLVENT: 539 g/L N-METHYL PYRROLIDONE

GROUP	C	FUNGICIDE
-------	----------	-----------

1 L

For control of a range of fungal diseases in turf as per the directions for use.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USE

(label code)

FRONT PAGE OF BOOKLET

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Chipco Baycor Fungicide

Active Constituent: 300 g/L BITERTANOL

Solvent: 539 g/L N-METHYL PYRROLIDONE

For control of a range of fungal diseases in turf as per the directions for use.

Storage and Disposal

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Dispose of at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Safety Directions

Harmful if swallowed. Will damage eyes. May irritate the skin. When using the product wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length butyl rubber gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 131126). If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

Material Safety Data Sheet

Additional information is listed in the Material Safety Data Sheet

Exclusion of Liability

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

NRA Approval No.: 56589/0403

IMPORTANT: READ THIS BOOKLET BEFORE USE

Ancillary panel attached to container

Storage and Disposal

Store in the closed, original container in a cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Dispose of at a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

Safety Directions

Harmful if swallowed. Will damage eyes. May irritate the skin. When using the product wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length butyl rubber gloves and goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone 131126). If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

Material Safety Data Sheet

Additional information is listed in the Material Safety Data Sheet

NRA Approval No.: 56589/0403

Chipco[®] and Baycor[®] are Registered Trademarks of Bayer.

Bayer Environmental Science
Bayer CropScience Pty Ltd
A.B.N. 87 000 226 022
391-393 Tooronga Rd
East Hawthorn Vic 3123
Phone: 03 9248-6888
Fax: 03 9248-6800
Website: www.bayercropscience.com.au

FOR 24 HOUR SPECIALIST ADVICE
IN EMERGENCY ONLY
PHONE **1800 033 111**

*Batch Number:

*Date of Manufacture:

*Bar code:

Label code

Bayer Environmental Science
A Business Group of Bayer CropScience

BOOKLET

DIRECTIONS FOR USE

Crop	Disease	Rate	Critical Comments
Kikuyu turf (NSW ONLY)	Helminthosporium leaf spot	2.5 L/ha	Severe outbreaks of Helminthosporium are favoured by warm, humid weather. Spray before the disease begins to restrict leaf and runner development. A repeat spray after two weeks may be necessary.
Turf (TAS ONLY)			
Couch turf	Black Helminthosporium (suppression of white Helminthosporium symptoms) (QLD NSW ONLY)	25-50 mL/100 m ²	Use high volume equipment. Low volume may be used provided turf is irrigated thoroughly after spraying. Apply during period of turf growth on a monthly program. Use higher rate during conditions of severe disease infection.
Turf	Dollar spot (QLD NSW VIC SA WA ONLY)	50 mL/100 m ²	Apply thoroughly on a monthly preventive spray program during the warmer months. Allow treatment to dry before turf is irrigated. Not for curative control.
	Spring dead spot (QLD NSW WA ONLY)		Tank mix with thiram 800 g/kg WP at 150 g/100 m ² plus soil wetting agent at 75 mL/100 m ² in a convenient volume of water (high or low volume). Make 6 applications at monthly intervals, starting early March (WA) or February (NSW). Apply to leaves that are wet, then water immediately after spraying.
	Winter Fusarium (VIC TAS ONLY)		Use high volume equipment. Low volume may be used provided turf is irrigated thoroughly after spraying. Preventive: apply thoroughly on a monthly program from end of March. Curative: if lesions are present, apply thoroughly and repeat at monthly intervals.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS**Fungicide Resistance Warning**

GROUP

C

FUNGICIDE

Chipco Baycor is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group C fungicide. Some naturally occurring individual fungi resistant to the product and other Group C fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group C fungicides, thus resulting in a reduction in efficacy. Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that result from failure of this product to control resistant fungi.

Mixing: Mix the required quantity of Chipco Baycor with water in the spray vat while stirring or with agitators operating.

BOOKLET (contd)

Compatibility: Chipco Baycor is compatible with Chipco Baythroid® Insecticide. The compatibilities given are based on world wide experience and are considered correct. However as changes in climatic conditions can alter the sensitivity of plants to mixtures of sprays we accept no responsibility for the behaviour of such mixtures.

Protection of Livestock, Wildlife, Fish, Crustaceans and Environment

Dangerous to fish. Do not contaminate ponds, waterways and drains with this product or used container. Do not feed grass clippings from treated areas to poultry or other animals.